5.50”

7.625”

7.875”

ORTHO®

WEED B GON® MAX®

FOR **SOUTHERN LAWNS** PARA CESPEDES DEL SUR

READY-TO-USE

KILLS WEEDS, NOT LAWNS™

KEEP OUT OF REACH OF CHILDREN

CAUTION See back panel booklet for additional precautionary statements.

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

PRECAUCION Consulte el folleto en el panel posterior para conocer los avisos de precaución adicionales.

Active Ingredients:

2,4-D, 2-ethylhexyl ester	0.184%
Mecoprop-p acid	0.022%
Dicamba acid	0.012%
Carfentrazone-ethyl	0.007%
Other Ingredients:	99.775%
TOTAL	100.000%

THIS PRODUCT CONTAINS: 2,4-Dichlorophenoxyacetic acid equivalent 0.122%. Isomer Specific by AOAC Methods.

FOR HOME LAWNS ONLY

OUTDOOR USE ONLY

FPO

LB2676

NET CONTENTS/CONTENIDO NETO 1 gal/3.78 l

5.75”

1 Gallon ORTHO® Pull ’N Spray®

EPA Reg No. 2217-905-239 (1gal)
Ortho Weed B Gon Max For Southern Lawns Ready -To-Use

EPA Reg No. 2217-905-239 (1gal)

Ortho Weed B Gon Max For Southern Lawns Ready -To-Use

4.3125”

4.50”

7.25”

7.625”

RIESGOS AMBIENTALES:

Este pesticida es tóxico para los peces y los invertebrados acuáticos y puede afectar de manera adversa a las plantas que no son el objetivo del tratamiento. Para proteger el medio ambiente, no permita que el pesticida se filtre o se introduzca en drenajes de tormenta, zanjas de drenaje, canalones o agua en superficie. Aplique este producto cuando el clima está tranquilo y cuando no hay pronóstico de lluvia para las siguientes 24 horas para ayudar a asegurar que el viento o la lluvia no sople o lave el pesticida fuera del área tratada.

Este producto químico tiene propiedades y características asociadas con las sustancias químicas detectadas en aguas subterráneas. La utilización de este producto químico en áreas donde los suelos son permeables, en particular donde el manto freático es superficial, puede derivar en la contaminación de las aguas subterráneas. La aplicación en torno a cisternas o pozos puede provocar la contaminación del agua potable o de las aguas subterráneas.

LA GARANTIA ORTHO®: Si por cualquier razón usted, el consumidor, no está satisfecho con este producto, envíenos por correo el comprobante original de compra para recibir un reembolso total del precio de compra.

ORTHO® Y SU ENTORNO

Su jardín es un lugar en el que su familia y sus mascotas disfrutan. Por este motivo, los productos Ortho® están diseñados cuidadosamente para proporcionar soluciones efectivas a los problemas de malezas. Para lograr los mejores resultados, siga las instrucciones de uso, almacenamiento y eliminación.

NOTA: En la medida que lo permita la ley aplicable, el comprador asume todos los riesgos de uso, almacenamiento o manipulación de este producto en caso de no utilizarlo de acuerdo con las instrucciones de la etiqueta.

8

ORTHO®

WEED B GON® MAX®

FOR **SOUTHERN LAWNS**
READY-TO-USE

PRODUCT FACTS Container treats more than 1000 weeds

Kills major listed broadleaf weeds including Dandelion, Clover, Oxalis, Chickweed, Knotweed, Plantain, Henbit, Spurge and many others.

Before

After

WHERE TO USE:

ON LAWNS – bahiagrass, bentgrass, bermudagrass (hybrid and common), buffalograss, centipedegrass, fescue, Kentucky bluegrass, perennial ryegrass, St. Augustinegrass (except ‘Floratam’) and zoysiagrass without harming lawn grasses when used as directed.

Do not allow people or pets to enter the treated area until spray has dried.

The ORTHO® Guarantee: If for any reason you, the consumer, are not satisfied with this product, mail us your original proof of purchase to obtain a full refund of your purchase price.

OPEN
Resealable Label for Directions & Precautions
Español ver página 5

F.P.O. 100%
0 71549 03946 2
1005

Questions, Comments or Medical Information
Preguntas, comentarios o información médica
1-800-225-2883 **www.ortho.com**

©2013 The ORTHO Group. World rights reserved.
Sold by **The ORTHO Group**
P.O. Box 190, Marysville, OH 43040

EPA Reg. No. 2217-905-239
EPA Est. No. 239-IA-3, 239-MS-001M, 58996-MO-1A, 085652-OH-1W
Superscript is first letter of lot number.

FPO

LB2893

4.75”

4.75”

EPA Reg No. 2217-905-239 (1gal)
Ortho Weed B Gon Max For Southern Lawns Ready -To-Use

4.3125”

4.3125”

7.25”

7.625”

STOP **READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Precautions and Restrictions

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.
- Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.
- Do not apply when windy.
- To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

Do not allow people or pets to enter the treated area until spray has dried.

PULL ‘N SPRAY® APPLICATOR

	• Remove sprayer from side carrier and unwrap hose completely. • Insert red plug at end of hose into red spout on cap **until it clicks**. • Flip up spout.
	• Twist nozzle at end of sprayer to adjust spray pattern.
	• Point sprayer away from body. • Grasp sprayer by the handle. • Slowly pull ring at bottom of sprayer handle until it stops to ready the sprayer.
	• Press and hold button on sprayer to begin spraying. • Pull ring at sprayer bottom again as needed to continue spraying.

HOW TO USE

Follow illustrations and instructions above. Twist nozzle at end of sprayer to adjust spray pattern. Point sprayer away from body and press sprayer button to begin spraying. Pull ring at sprayer bottom again as needed to continue spraying. Aim at center of weed and spray lightly to cover.

WHEN TO APPLY

- Spray when weeds are actively growing.
- Spray when air temperatures are below 90°F and are not expected to exceed 90°F for 6 hours.
- Hard to kill weeds may require a repeat application in 30 days.
- Rain or watering 6 hours after application will not wash away effectiveness.
- Weeds in newly seeded lawns may be sprayed after the lawn has been mowed 3 times.
- The number of spot treatments is limited to 2 per year.

IMPORTANT

Do not use this product for controlling weeds in vegetable gardens, flower beds or around shrubs or ornamental plants. Do not use on or around fruits, vegetables or flowers. Temporary discoloration may occur on St. Augustinegrass, centipedegrass, and bentgrass lawns. Do not apply as a fine mist because of potential for injury to desirable plants.

BROADLEAF WEEDS CONTROLLED

Ortho® Weed B Gon® MAX® for Southern Lawns Ready-to-Use will control or suppress the following broadleaf weeds.

Amaranth (purple), Amaranth (slender), Aster (heath), Aster (white heath), Aster (purple), Aster (white prairie), Artichoke, Austrian fieldcress, Bedstraw (catch-weed), Bedstraw (smooth), Bedstraw (northern), Bedstraw (pineywoods), Beggar-ticks (hairy), Betony (Florida), Bindweed (field), Bindweed (hedge), Bitter Wintercress, Bitter-cress (hairy), Bitterweed, Black-eyed Susan, Black medic, Blood-flower milkweed, Blue lettuce, Brassbuttons, Bristly oxtongue, Broomweed, Buckhorn, Bull-nettle, Bur-clover, Burdock, Burweed (lawn), Buttercup (bulbous), Buttercup (creeping), Buttonweed (common [poorjoe]), Buttonweed (Virginia), Carolina geranium, Carpetweed, Catnip, Catsear (spotted [false dandelion]), Chamber bitter, Chickweed (common), Chickweed (mouseear), Chickweed (sticky), Chicory, Cinquefoil, Clover (crimson), Clover (Alyce), Clover (hop), Clover (rabbit-foot), Clover (red), Clover (strawberry), Clover (yellow sweet), Clover (white sweet), Clover (white), Cockle, Cocklebur, Cocklebur (spiny), Common mullein, Corn spurry, Creeping beggar-weed, Creeping Jenny (creeping Charlie), Cupid’s shaving brush, Cudweed, Daisy (English), Daisy (oxeye), Dandelion (common), Dandelion (false), Day-flower, Deadnettle, Dichondra, Dock (broadleaf), Dock (curly), Dogbane, Dog-fennel, Dollarweed (Pennywort), Dove-weed, Elderberry, Falseflax, False sunflower, Fiddleneck, Filaree (redstem), Filaree (whitestem), Fleabane (daisy [annual]), Fleabane (rough), Galinsoga (hairy), Galinsoga (smallflower), Goldenrod, Ground ivy, Groundsel, Gumweed, Hairy fleabane, Hawkweed (meadow), Hawkweed (mouse-ear), Hawkweed (orange), Hawkweed (yellow), Healall, Heart-leaf drymary, Hemp, Henbit, Hoary-cress, Horsenettle, Horseweed, Jimsonweed, Johnny-jumpup violet, Knawel, Knotweed (prostrate), Kochia, Lambsquarters, Lespedeza, Mallow (alkali), Mallow (bristly), Mallow (common), Mallow (dwarf), Mallow (Venice), Marcela, Matchweed, Mexican-weed, Moneywort, Morningglory (bigroot), Morningglory (ivyleaf), Morningglory (red), Morningglory (tall), Morningglory (woolly), Mugwort, Mustard (ball), Mustard (black), Mustard (blue), Mustard (hedge), Mustard (Indian), Mustard (tansy), Mustard (tumble), Mustard (white), Mustard (wormseed), Nettle (stinging), Nettle (tall), Nettle (wood), Parsley-piert, Parsnip, Pearlwort, Pennycress, Pepperweed (perennial), Pepperweed (Virginia [Peppergrass]), Pigweed (prostrate), Pigweed (redroot), Pigweed (Russian), Pigweed (smooth), Pigweed (tumble), Pineapple weed, Plains coreopsis (tickseed), Plantain (black-seed), Plantain (bracted), Plantain (broadleaf), Plantain (buckhorn), Plantain (hoary), Plantain (narrow-leaf), Plantain (slender), Plantain (woolly), Poison ivy, Poison oak, Pokeweed, Prairie sunflower, Prickly lettuce (compass plant), Prickly sida, Prostrate knotweed, Puncturevine, Purslane (common), Pusley (Brazilian), Pusley (Florida), Pusley (large flower), Ragweed (bur), Ragweed (common), Ragweed (lance-leaf), Ragweed (western), Redstem filaree, Rough cinquefoil, Scarlet pimpernel, Shepherd’s purse, Smartweed (ladysthumb), Smartweed (pale), Smartweed (Pennsylvania), Smooth chaff-flower, Smooth dock, Sorrel (red [sheep]), Sowthistle (annual), Sowthistle (spiny), Spanish needles, Speedwell (bi-lobed), Speedwell (common), Speedwell (corn), Speedwell (ivyleaf), Speedwell (Persian), Speedwell (purslane), Speedwell (slender), Speedwell (snow), Speedwell (thyme-leaf), Spiny amaranth, Spurge (spotted), Spurge (prostrate), Star of Bethlehem, Strawberry (Indian mock), Tansy ragwort, Tanweed, Thistle (blessed), Thistle (bull), Thistle (Canada), Thistle (Flodman’s), Thistle (musk), Thistle (plumeless), Thistle (prairie), Thistle (Russian), Thistle (Scotch), Thistle (wavy-leaf), Thistle (yellow), Trailing crown-vetch, Velvetleaf, Vervain (blue), Vervain (hoary), Vervain (prostrate), Vervain (tall), Vetch (bird), Vetch (hairy), Vetch (milk), Vetch (narrow-leaf), Vetch (wild), Virginia creeper, Western clematis, Western salsify, Wild aster, Wild buckweat, Wild carrot, Wild four-o-clock, Wild garlic, Wild lettuce, Wild marigold, Wild mustard, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild violet, Woodsorrel (creeping), Woodsorrel (violet), Woodsorrel (yellow [*Oxalis*]), Woolly croton, Wormseed, Yarrow (common), Yarrow (western), Yarrow (yellow), Yellow Rocket

4.75”

4.75”

EPA Reg No. 2217-905-239 (1gal)
Ortho Weed B Gon Max For Southern Lawns Ready -To-Use

4.3125”

4.3125”

7.25”

7.625”

STORAGE AND DISPOSAL

PESTICIDE STORAGE	Do not contaminate water, food, or feed by storage and disposal. Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Completely discharge product in sprayer prior to storage. Twist sprayer nozzle to the OFF position. Flip down spout on cap. NO NEED TO DISCONNECT SPRAYER HOSE FROM CAP. Place sprayer back inside carrier on bottle with the nozzle facing down.
PESTICIDE DISPOSAL AND CONTAINER HANDLING	Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instruction. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

IF ON SKIN	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-225-2883 for emergency medical treatment advice.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

THE ORTHO® GUARANTEE: If for any reason you, the consumer, are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

ORTHO® & YOUR ENVIRONMENT

Your yard is a place for your family and pets to enjoy. That’s why Ortho® products are designed with care to provide effective solutions to weed problems. Follow instructions for appropriate usage, storage and disposal.

NOTICE: To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.

ORTHO® Weed B Gon® MAX®
Para Céspedes del Sur
Listo para Usar

ESPECIFICACIONES DEL PRODUCTO
El envase elimina más de 1000 malezas

Elimina las principales malezas de hoja ancha que se enumeran, como diente de león, trébol, oxalis, álsine, sanguinaria, plántago, ortiga mansa, lechetrezna y muchas otras.

DONDE DEBE UTILIZAR EL PRODUCTO	**SOBRE CESPEDES** – césped bahía, agrostis, césped Bermuda (híbrido y común), césped búfalo, césped ciempiés, festuca, hierba azul de Kentucky, raigrás perenne, césped San Agustín (excepto ‘Floratam’) y césped zoysia; no daña el césped si se utiliza según las instrucciones.

PARE **LEA PRIMERO LA ETIQUETA COMPLETA. OBSERVE TODAS LAS PRECAUCIONES Y SIGA LAS INSTRUCCIONES DETENIDAMENTE.**

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.

Restricciones y precauciones

- No aplique este producto de forma que entre en contacto con otras personas o mascotas, ya sea directamente o a través de la desviación del producto. Mantenga a las personas y las mascotas lejos del área durante la aplicación del producto.
- No aplique el producto directamente al o cerca del agua, drenajes de tormenta, canaletas, alcantarillas o diques de drenaje.
- No aplique el producto a menos de 8 metros de ríos, estanques de peces, lagos, riachuelos, embalses, pantanos, estuarios, bahías y océanos.
- No aplique el producto cuando hay viento.
- Para evitar el escurrimiento del producto, no riegue el área tratada en exceso hasta el punto de escurrimiento, ni lo aplique en condiciones de lluvia o en días con pronóstico de lluvia.

No permita que las personas o las mascotas ingresen al área tratada hasta que el rociado se haya secado.

EL APLICADOR PULL ‘N SPRAY®

	• Saque el rociador del cargador lateral y desenrolle completamente la manguera. • Inserte el tapón rojo al final de la manguera en la boquilla roja de la tapa **hasta que haga clic**. • Abra la boquilla hacia arriba.
	• Gire la boquilla en el extremo del rociador para ajustar el patrón de rocío.
	• Dirija el rociador lejos del cuerpo. • Agarre el rociador por la manija. • Lentamente jale el anillo en la parte inferior de la manija del rociador hasta que pare para que el rociador esté listo.
	• Presione y sostenga el botón en el rociador para empezar a rociar. • Jale nuevamente el anillo en la parte inferior del rociador conforme lo necesite para continuar rociando.

4.75”

4.75”

EPA Reg No. 2217-905-239 (1gal)
Ortho Weed B Gon Max For Southern Lawns Ready -To-Use

4.3125”

4.3125”

7.25”

7.625”

COMO USAR

Siga las ilustraciones e instrucciones que figuran arriba. Gire la boquilla en el extremo del rociador para ajustar el patrón de rociado. Dirija el rociador lejos de su cuerpo y oprima el botón del rociador para empezar a rociar. Tire nuevamente del anillo en la parte inferior del rociador, según sea necesario, para continuar rociando. Apunte al centro de la maleza y rocíe para cubrirla ligeramente.

CUANDO APLICAR

- Rocíe cuando las malezas estén creciendo activamente.
- Rocíe cuando la temperatura sea inferior a 32°C y no se espere que exceda los 32°C por 6 horas.
- Las malezas resistentes pueden requerir otra aplicación a los 30 días.
- La lluvia o el riego 6 horas después de la aplicación no reducirán su eficacia.
- Las malezas en céspedes recién sembrados se pueden rociar después de que se haya cortado el césped 3 veces.
- El número de tratamientos focalizados no debe exceder de 2 por año.

IMPORTANTE

No utilice este producto para controlar las malezas en hortalizas, canteros de flores o en torno a arbustos o plantas ornamentales. No lo utilice sobre o cerca de frutas, verduras o flores. Se puede producir una decoloración temporal en el césped San Agustín, césped ciempiés y agrostis. No aplique como rocío fino ya que puede dañar las plantas que no desea eliminar.

MALEZAS DE HOJA ANCHA CONTROLADAS

Ortho® Weed B Gon® MAX® para Céspedes del Sur Listo para Usar controlará o suprimirá las siguientes malezas de hoja ancha.

Amaranto (púrpura), Bledo blanco, Aster ericoides, Aster (brezo blanco), Aster púrpura, Aster de pradera blanca, Alcaucil, Rorippa austríaca, Cuajaleche (Galium verum), Cuajaleche (Galium mollugo), Cuajaleche del norte (Galium septentrionale), Cuajaleche (pineywoods), Amor seco (bidens pilosa), Betónica de Florida, Correhuela (silvestre), Correhuela mayor, Berro de altura, Mastuerzo amargo piloso, Cola de caballo, Jovinova, Alfalfa lupulina, Flor de sangre, Lechuga azul, Cotula, Raspalenguas, Escobilla, Llantén, Pringamosa, Trébol de carretilla, Bardana, Cadillo, Ranúnculo bulboso, Ranúnculo rastrero, Hierba de botón común (“poorjoe”), Hierba de botón de Virginia, Geranio de Carolina, Verdolaga alfombra, Hierba gatuna, Hierba del halcón veteada (falso diente de león), Filantus, Alsine (común), Alsine (oreja de ratón), Alsine (pegajosa), Achicoria, Potentilla, Trébol carmesí, Trébol de una hoja, Trébol campestre, Trébol pata de conejo, Trébol rojo, Trébol frutilla, Trébol amarillo dulce, Trébol blanco dulce, Trébol blanco, Neguilla, Bardana menor, Xanthium Spinosum, Gordolobo común, Espérgula, Trébol español de Kaimi, Lysimachia nummularia (“Creeping Jenny”, “Creeping Charlie”), Botón rosado (“Cupid’s shaving brush”), Siempreviva, Margarita inglesa, Margarita mayor, Diente de león común, Falso diente de león, Comelina, Ortiga blanca, Oreja de ratón, Lengua de vaca, Romaza, Flamenquilla, Manzanilla cimarrona, Acaricoba (“Pennywort”), Croton Setigerus, Baya del saúco, Falso lino, Falso girasol, Amsinckia (“Fiddleneck”), Alfilerillo de pastor, Hierba almizclera, Erigeron (margarita [anual]), Erigeron áspero, Galinsoga cuadriradiata, Galinsoga parviflora, Vara de oro, Hiedra terrestre, Hierba cana, Grindelia, Hierba del naciente, Hieracium caespitosum, Hieracium pilocella, Hieracium aurantiacum, Hieracium fendleri, Hierba mora, Drimaria (hoja de corazón), Cáñamo, Ortiga mansa, Draba, Solanum carolinense, Erigeron canadensis, Estramonio, Viola tricolor, Scleranthus annuus, Cien nudos, Pinillo, Quinoa, Lespedeza, Malva leprosa, Modiola caroliniana, Malva común, Malva neglecta, Hibiscus trionum, Marcela, Lippia, Maleza mexicana, Hierba de la moneda, Ipomoea de raíz grande, Ipomoea (hoja de hiedra), Ipomoea roja, Ipomoea purpúrea, Ipomoea lanosa, Artemisia, Tamarillas, Mostaza negra, Mostaza azul, Erísimo, Mostaza india, Descurainia pinnata, Nabo chileno, Mostaza blanca, Erysimum cheiranthoides, Ortiga, Ortiga delgada, Laportea canadensis, Alquimila arvense, Pastinaca, Sagina subulata, Carraspique, Lepidio perenne, Lepidio de Virginia (“Peppergrass”), Bledo postrado, Yuyo de raíces coloradas, Cardo ruso, Bledo, Bledo blanco, Piñas de algas, Coreopsis tinctoria (dalia de mar), Llantén de semilla negra, Plántago patagónico, Llantén mayor, Llantén menor, Llantén medio, Plántago menor, Plántago heterophylla, Hierba del pastor, Hiedra venenosa, Roble venenoso, Paloma baya, Girasol de la pradera, Lechuga silvestre (“compass plant”), Sida spinosa, Sanguinaria, Roseta francesa, Verdolaga, Yerba del sapo, Ipecacuana blanca, Ipecacuana de flores grandes, Falsa ambrosía, Ambrosía común, Ambrosía bidentata, Ambrosía americana, Alfilerillo, Potentilla norvegica, Pimpinela escarlata, Bolsa del pastor, Persicaria, Persicaria lapathifolia, Persicaria pensylvánica, Alternanthera paronychioides, Rumex altissimus, Amargosa, Cerraja, Cerraja brava, Bidente, Verónica biloba, Verónica, Verónica arvensis, Hiedrezuela terrestre, Verónica pérsica, Verónica peregrina, Verónica filiformis, Verónica nivea, Verónica oltensis, Amaranto espinoso, Yerba de la golondrina, Bledo rastrero, Estrella de Belén, Fresa de la India, Hierba de Santiago, Polígono de pantano, Cardo bendito, Cardo negro, Cardo de Canadá, Cardo Flodman, Cardo pendiente, Cardo, Cardo lámina, Cardo ruso, Cardo escocés, Cardo (Cirsium undulatum), Cardo amarillo, Coronilla varia, Abutilón, Verbena azul, Verbena stricta, Verbena bracteata, Verbena bonariensis, Veza de pájaro, Veza vellosa, Tragacanto, Vicia angustifolia, Veza silvestre, Enamorada del muro, Clemátide, Clavelón, Aster silvestre, Trigo sarraceno silvestre, Zanahoria silvestre, Maravilla, Ajo silvestre, Lechuga silvestre, Chinchilla, Mostaza silvestre, Cebolla silvestre, Pastinaca silvestre, Rabanillo, Rábano silvestre, Frutilla silvestre, Batata silvestre, Violeta silvestre, Vinagrillo, Oxalis violácea, Oxalis stricta (Oxalis), Hierba de cabra, Epazote, Milenrama común, Milenrama occidental, Milenrama amarilla, Jaramago

ALMACENAMIENTO Y ELIMINACION

ALMACENAMIENTO DEL PESTICIDA	No contamine el agua, la comida o el alimento para animales, al almacenarlo o eliminarlo. Evite que el producto se congele. Mantenga el pesticida en su envase original. No lo coloque en envases de comidas o bebidas. Evite la contaminación de comestibles y alimentos para animales. Almacene el producto en un lugar fresco y seco, preferentemente en un área de almacenamiento bajo llave. Descargue todo el producto en el rociador antes de almacenarlo. Gire la boquilla del rociador a la posición de OFF (APAGADO). Baje la boquilla en la tapa. NO ES NECESARIO DESCONECTAR LA MANGUERA DEL ROCIADOR DE LA TAPA. Vuelva a colocar el rociador dentro del soporte de la botella con la boquilla hacia abajo.
ELIMINACION DEL PESTICIDA Y MANEJO DEL ENVASE	Envase no rellenable. No vuelva a utilizar o llenar este envase. **Si está vacío:** Colóquelo en la basura o, si corresponde, ofrézcalo para reciclaje. **Si está parcialmente lleno:** Llame a su agencia de desechos sólidos para obtener instrucciones sobre cómo eliminar el producto. Nunca elimine el producto no utilizado en drenajes internos o externos.

AVISOS DE PRECAUCION

RIESGOS PARA SERES HUMANOS Y ANIMALES DOMESTICOS

PRECAUCION: Este producto es nocivo si se absorbe por la piel. Evite el contacto con la piel, los ojos o la ropa. Lávese bien con agua y jabón después de manipular el producto y antes de comer, beber, masticar chicle, usar tabaco o usar el baño.

PRIMEROS AUXILIOS

SI EL PRODUCTO ENTRA EN CONTACTO CON LA PIEL	• Quítese la ropa contaminada. • Lave inmediatamente la piel con abundante agua durante 15 a 20 minutos. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. También puede llamar al 1-800-225-2883 para obtener asesoramiento sobre el tratamiento médico de emergencia.

4.75”

4.75”

EPA Reg No. 2217-905-239 (1gal)
Ortho Weed B Gon Max For Southern Lawns Ready -To-Use

4.375”

7.375”

7.625”

STORAGE AND DISPOSAL

PESTICIDE STORAGE	Do not contaminate water, food, or feed by storage and disposal. Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Completely discharge product in sprayer prior to storage. Twist sprayer nozzle to the OFF position. Flip down spout on cap. NO NEED TO DISCONNECT SPRAYER HOSE FROM CAP. Place sprayer back inside carrier on bottle with the nozzle facing down.
PESTICIDE DISPOSAL AND CONTAINER HANDLING	Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instruction. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

IF ON SKIN	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-225-2883 for emergency medical treatment advice.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

THE ORTHO® GUARANTEE: If for any reason you, the consumer, are not satisfied with this product, mail us your original proof of purchase to obtain a full refund of your purchase price.

NOTICE: To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.

Questions, Comments or Medical Information
Preguntas, comentarios o información médica
call 1-800-225-2883 www.ortho.com

©2013 The ORTHO Group. World rights reserved.
Sold by **The ORTHO Group**
P.O. Box 190, Marysville, OH 43040

EPA Reg. No. 2217-905-239
EPA Est. No. 239-IA-3I, 239-MS-001M,
58996-MO-1A, 085652-OH-1W
Superscript is first letter of lot number. LB2893

PRESS TO RESEAL PRESIONE PARA VOLVER A CERRAR

5.50”

EPA Reg No. 2217-905-239 (1gal)
Ortho Weed B Gon Max For Southern Lawns Ready -To-Use